



GmA	Goldsboro sandy loam, 0 to 2 percent slopes	37.70	3.1%		llw					125		700						
LMA	Lucy loamy sand, 0 to 2 percent slopes	36.67	3.0%		lls													
TuA	Tifton sandy loam, 0 to 2 percent slopes	35.71	2.9%		lw													
LtA	Lynchburg sandy loam, 0 to 2 percent slopes	31.83	2.6%		llw													
WeB	Wagram loamy sand, 2 to 5 percent slopes	28.59	2.3%		lls													
RgA	Red Bay loamy sand, 0 to 2 percent slopes	26.17	2.1%		l													
OeB2	Orangeburg loamy sand, 2 to 5 percent slopes, moderately eroded	24.54	2.0%		lle													
AdA	Albany sand, 0 to 2 percent slopes	15.68	1.3%		lllw													
LpB	Lakeland sand, 0 to 5 percent slopes	15.58	1.3%		llls	llls		7		55								7
IgA	Irvington sandy loam, 0 to 2 percent slopes	15.17	1.2%		llw													
WeA	Wagram loamy sand, 0 to 2 percent slopes	11.58	0.9%		lls													
SUB2	Sawyer-Susquehanna loamy sands, 2 to 5 percent slopes, eroded	9.93	0.8%		lVe													
CoC2	Carnegie sandy loam, 5 to 8 percent slopes, moderately eroded	7.39	0.6%		lVe													
CoC2	Cuthbert-Orangeburg complex, 5 to 8 percent slopes	3.90	0.3%		lVe													
CoB2	Carnegie sandy loam, 2 to 5 percent slopes, moderately eroded	2.44	0.2%		llle													
RgB2	Red Bay loamy sand, 2 to 5 percent slopes, eroded	1.58	0.1%		llle													
W	Water	1.44	0.1%															
Weighted Average							1.3	3.1	1.1	45.5	24.6	357.9	142.4	22.7	0.8	3.8		

*c: Using Capabilities Class Dominant Condition Aggregation Method